

Answers to Study Questions - Topic 18

1. (a) \bar{T} is the temporal average of T :

$$\begin{aligned}\bar{T} &= \frac{1}{6} \sum_{t=0}^5 T(t) \\ &= \frac{1}{6} (12.6 + 11.2 + 11.9 + 13.1 + 12.0 + 11.8) \\ &= \underline{12.1^\circ\text{C}}\end{aligned}$$

(b) T' is the deviation of a single measured value from the temporal average of the time series (\bar{T}). For the time step at 40 min we can write:

$$\begin{aligned}T(40\text{min}) &= \bar{T} + T'(40\text{min}) \\ T'(40\text{min}) &= T(40\text{min}) - \bar{T} \\ &= 13.1^\circ\text{C} - 12.1^\circ\text{C} \\ &= \underline{1\text{ K}}\end{aligned}$$

(c) T'^2 is the squared deviation for a single measured value from the temporal average of the time series. For the time step at 20 min we can write: (\bar{T}):

$$\begin{aligned}T'^2(20\text{min}) &= (T(20\text{min}) - \bar{T})^2 \\ &= (11.2^\circ\text{C} - 12.1^\circ\text{C})^2 \\ &= (-0.9^\circ\text{C})^2 \\ &= \underline{0.81\text{ K}^2}\end{aligned}$$

(d) \bar{T}' is the average deviation in the time series. By definition, this term must be always zero, i.e $\bar{T}' = 0$ (or more general $\bar{a}' = 0$, where a

is any parameter or term). Let's check this:

$$\begin{aligned}
\overline{T'} &= \frac{1}{6} \sum_{t=0}^5 (T(t) - \overline{T}) \\
&= \frac{1}{6} ((12.6 - 12.1) + (11.2 - 12.1) + (11.9 - 12.1) \\
&\quad + (13.1 - 12.1) + (12.0 - 12.1) + (11.8 - 12.1)) \\
&= \frac{1}{6} (0.5 - 0.9 - 0.2 + 1.0 - 0.1 - 0.3) \\
&= \underline{0 \text{ K}}
\end{aligned}$$

(e) $\overline{T'^2}$ is the the variance of the time series of T . It is defined as the average of the squared deviations of all time steps. This term is not necessarily zero¹

$$\begin{aligned}
\overline{T'^2} &= \frac{1}{6} \sum_{t=0}^5 (T(t) - \overline{T})^2 \\
&= \frac{1}{6} \left((12.6 - 12.1)^2 + (11.2 - 12.1)^2 + (11.9 - 12.1)^2 \right. \\
&\quad \left. + (13.1 - 12.1)^2 + (12.0 - 12.1)^2 + (11.8 - 12.1)^2 \right) \\
&= \frac{1}{6} (0.25 + 0.81 + 0.04 + 1 + 0.01 + 0.09) \\
&= \underline{0.37 \text{ K}^2}
\end{aligned}$$

(f) $\overline{T'^2}$ is the temporal average of the deviations squared. Note the fine (but essential) difference of the overbar that does not include the square, and because $\overline{T'} = 0$ (see (d)) also its square is zero:

$$\overline{T'^2} = \overline{T'} \times \overline{T'}$$

¹Note for those who have taken statistics: You might be used to the *unbiased variance*, where you divide by $N - 1$, not N :

$$\overline{T'^2} = \frac{1}{N-1} \sum_{t=0}^{N-1} (T(t) - \overline{T})^2 = \frac{1}{5} \sum_{t=0}^5 (T(t) - \overline{T})^2$$

However, in atmospheric turbulence studies, we prefer the biased variance (divided by N , as shown in the question above and the lecture slides where we divide by N). The biased variance is a good estimation of the dispersion of a sample of observations, but not necessarily the best measure of the whole population of possible observations.

$$\begin{aligned}
&= 0 \text{ K} \times 0 \text{ K} \\
&= \underline{0 \text{ K}^2}
\end{aligned}$$

2. (a) The average of a constant is equal the constant itself

$$\bar{5} = \frac{1}{1} \sum_{i=0}^0 5 = 5$$

(b) The average of a constant times a value is equal the constant times the averaged value:

$$\overline{8v} = 8 \times \bar{v}$$

(c) Similarly we can regard temporal averages as constants (they are not changing over time):

$$\overline{\overline{Tp}} = \overline{T}\overline{p}$$

(d) The average of an averaged value is equal the averaged value

$$\overline{\bar{u}} = \bar{u}$$

(e) Similar to 1(d):

$$\overline{q'} = 0$$

(f) Again, the first term is zero - so is it's product:

$$\overline{T'} \times \overline{w} = 0 \times \overline{w} = 0$$

(g)

$$\overline{3q'} = 3 \times \overline{q'} = 3 \times 0 = 0$$

(h)

$$\overline{w' \times \bar{u}} = \overline{w'} \times \bar{u} = 0 \times \bar{u} = 0$$

(i) Combining the results from (c) and (e):

$$\begin{aligned}
\overline{\overline{Tp}} &= \overline{T} \times \overline{(\bar{p} + p')} \\
&= \overline{T}\overline{p} + \overline{T} \times \underbrace{\overline{p'}}_{=0} \\
&= \overline{T}\overline{p}
\end{aligned}$$

(j) In a first step apply Reynold's decomposition to both, w and T . One of the resulting terms is a covariance ($\overline{w'T'}$) which is not zero (see lecture):

$$\begin{aligned}
 \overline{wT} &= \overline{(\bar{w} + w') \times (\bar{T} + T')} \\
 &= \overline{\bar{w}\bar{T}} + \underbrace{\overline{wT'}}_{=0} + \underbrace{\overline{w'\bar{T}}}_{=0} + \overline{w'T'} \\
 &= \overline{w}\bar{T} + \underbrace{\overline{w'T'}}_{\text{Covariance}}
 \end{aligned}$$

3. Calculate the following parameters if $\bar{u} = 4 \text{ m s}^{-1}$, $\bar{v} = 0 \text{ m s}^{-1}$, $\bar{w} = 0 \text{ m s}^{-1}$, $\sigma_u = 0.4 \text{ m s}^{-1}$, $\sigma_v = 0.2 \text{ m s}^{-1}$, and $\sigma_w = 0.1 \text{ m s}^{-1}$.

(a) I_u is the turbulence intensity of the longitudinal wind component u . I_u has no unit.

$$\begin{aligned}
 I_u &= \sigma_u / \bar{u} \\
 &= 0.4 \text{ m s}^{-1} / 4 \text{ m s}^{-1} \\
 &= \underline{0.1}
 \end{aligned}$$

(b) I_w is the turbulence intensity of the vertical wind component w . I_w has no unit.

$$\begin{aligned}
 I_w &= \sigma_w / \bar{u} \\
 &= 0.1 \text{ m s}^{-1} / 4 \text{ m s}^{-1} \\
 &= \underline{0.025}
 \end{aligned}$$

(c) $\overline{w'^2}$ is the variance of the vertical wind component w . The variance is the square of the standard deviation σ_w :

$$\begin{aligned}
 \overline{w'^2} &= \sigma_w^2 \\
 &= (0.1 \text{ m s}^{-1})^2 \\
 &= \underline{0.01 \text{ m}^2 \text{ s}^{-2}}
 \end{aligned}$$

(d)

$$\begin{aligned}
 \overline{u'^2} + \overline{v'^2} + \overline{w'^2} &= \sigma_u^2 + \sigma_v^2 + \sigma_w^2 \\
 &= (0.4 \text{ m s}^{-1})^2 + (0.2 \text{ m s}^{-1})^2 + (0.1 \text{ m s}^{-1})^2 \\
 &= 0.16 \text{ m}^2 \text{ s}^{-2} + 0.04 \text{ m}^2 \text{ s}^{-2} + 0.01 \text{ m}^2 \text{ s}^{-2} \\
 &= \underline{0.21 \text{ m}^2 \text{ s}^{-2}}
 \end{aligned}$$

(e) MKE/m is the mean kinetic energy per unit mass (lecture 18, slide 15), i.e.

$$\begin{aligned}
 MKE/m &= \frac{1}{2}(\bar{u}^2 + \bar{v}^2 + \bar{w}^2) \\
 &= \frac{1}{2}(4^2 + 0^2 + 0^2) \\
 &= \frac{1}{2}(16) \\
 &= \underline{8 \text{ m}^2 \text{ s}^{-2}}
 \end{aligned}$$

(f) \bar{e} is the turbulent kinetic energy per unit mass (lecture 18, slide 15) i.e.

$$\begin{aligned}
 \bar{e} &= \frac{1}{2}(\bar{u'^2} + \bar{v'^2} + \bar{w'^2}) \\
 &= \frac{1}{2}(\sigma_u^2 + \sigma_v^2 + \sigma_w^2) \\
 &= \frac{1}{2} \left((0.4 \text{ m s}^{-1})^2 + (0.2 \text{ m s}^{-1})^2 + (0.1 \text{ m s}^{-1})^2 \right) \\
 &= \frac{1}{2}(0.16 \text{ m}^2 \text{ s}^{-2} + 0.04 \text{ m}^2 \text{ s}^{-2} + 0.01 \text{ m}^2 \text{ s}^{-2}) \\
 &= \underline{0.105 \text{ m}^2 \text{ s}^{-2}}
 \end{aligned}$$